

**Medication Guide**  
**Oxycodone (ox-ee-CO-dohn) Hydrochloride**  
**Oral Solution, CII**

**Oxycodone Hydrochloride Oral Solution is:**

- A strong prescription pain medicine that contains an opioid (narcotic) that is used to manage pain severe enough to require an opioid pain medicine, when other pain treatments such as non-opioid pain medicines do not treat your pain well enough or you cannot tolerate them.
- An opioid pain medicine that can put you at risk for overdose and death. Even if you take your dose correctly as prescribed, you are at risk for opioid addiction, abuse, and misuse that can lead to death.

**Important information about Oxycodone Hydrochloride Oral Solution:**

- **Get emergency help right away if you take too much Oxycodone Hydrochloride Oral Solution (overdose).** When you first start taking Oxycodone Hydrochloride Oral Solution, when your dose is changed, or if you take too much (overdose), serious or life-threatening breathing problems that can lead to death may occur.
- Taking Oxycodone Hydrochloride Oral Solution with other opioid medicines, benzodiazepines, alcohol, or other central nervous system depressants (including street drugs) can cause severe drowsiness, decreased awareness, breathing problems, coma, and death.
- Never give anyone else your Oxycodone Hydrochloride Oral Solution. They could die from taking it. Selling or giving away Oxycodone Hydrochloride Oral Solution is against the law.
- Store Oxycodone Hydrochloride Oral Solution securely, out of sight and reach of children, and in a location not accessible by others, including visitors to the home.

**Do not take Oxycodone Hydrochloride Oral Solution if you have:**

- severe asthma, trouble breathing, or other lung problems.
- a bowel blockage or have narrowing of the stomach or intestines.
- an allergy to oxycodone or any of the ingredients in Oxycodone Hydrochloride Oral Solution.

**Before taking Oxycodone Hydrochloride Oral Solution, tell your healthcare provider if you have a history of:**

- head injury, seizures
- liver, kidney, thyroid problems
- problems urinating
- pancreas or gallbladder problems
- abuse of street or prescription drugs, alcohol addiction, or mental health problems.

**Tell your healthcare provider if you are:**

- **pregnant or planning to become pregnant.** Prolonged use of Oxycodone Hydrochloride Oral Solution during pregnancy can cause withdrawal symptoms in your newborn baby that could be life-threatening if not recognized and treated.
- **breastfeeding.** Oxycodone Hydrochloride Oral Solution passes into breast milk and may harm your baby.
- taking prescription or over-the-counter medicines, vitamins, or herbal supplements. Taking Oxycodone Hydrochloride Oral Solution with certain other medicines can cause serious side effects that could lead to death.

**When taking Oxycodone Hydrochloride Oral Solution:**

- Do not change your dose. Take Oxycodone Hydrochloride Oral Solution exactly as prescribed by your healthcare provider. Use the lowest dose possible for the shortest time needed.
- Always use the syringe that comes with Oxycodone Hydrochloride Oral Solution to correctly measure your dose. Never use a household teaspoon or tablespoon to measure Oxycodone Hydrochloride Oral Solution.
- Take your prescribed dose every 4-6 hours. Do not take more than your prescribed dose. If you miss a dose, take your next dose at your usual time.
- Call your healthcare provider if the dose you are taking does not control your pain.
- If you have been taking Oxycodone Hydrochloride Oral Solution regularly, do not stop taking Oxycodone Hydrochloride Oral Solution without talking to your healthcare provider.
- Dispose of expired, unwanted, or unused Oxycodone Hydrochloride Oral Solution by promptly flushing down the toilet, if a drug take-back option is not readily available. Visit [www.fda.gov/drugdisposal](http://www.fda.gov/drugdisposal) for additional information on disposal of unused medicines.

**While taking Oxycodone Hydrochloride Oral Solution DO NOT:**

- Drive or operate heavy machinery, until you know how Oxycodone Hydrochloride Oral Solution affects you. Oxycodone Hydrochloride Oral Solution can make you sleepy, dizzy, or lightheaded.
- Drink alcohol or use prescription or over-the-counter medicines that contain alcohol. Using products containing alcohol during treatment with Oxycodone Hydrochloride Oral Solution may cause you to overdose and die.

**The possible side effects of Oxycodone Hydrochloride Oral Solution:**

- constipation, nausea, sleepiness, vomiting, tiredness, headache, dizziness, abdominal pain. Call your healthcare provider if you have any of these symptoms and they are severe.

**Get emergency medical help if you have:**

- trouble breathing, shortness of breath, fast heartbeat, chest pain, swelling of your face, tongue, or throat, extreme drowsiness, light-headedness when changing positions, feeling faint, agitation, high body temperature, trouble walking, stiff muscles, or mental changes such as confusion.

These are not all the possible side effects of Oxycodone Hydrochloride Oral Solution. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. **For more information, go to [dailymed.nlm.nih.gov](http://dailymed.nlm.nih.gov).**

Manufactured by: **VistaPharm, Inc., Largo, Florida**  
[www.vistapharm.com](http://www.vistapharm.com) or call 1-888-655-1505

**This Medication Guide has been approved by the U.S. Food and Drug Administration.**

**Manufactured by:**

***VistaPharm***<sup>®</sup>

Largo, FL 33771

VP 2100R3

04/19

Proof **(A)** B C D E F G

PLEASE SIGN AND RETURN

OK to Print

Send Another Proof

Signature \_\_\_\_\_ Date \_\_\_\_\_

Job No. 191038 Item No. VP2100R3 Rev. 04/19

Date Sent 5-02-19 Date Returned \_\_\_\_\_

Art Sent

Art Returned

This proof is printed on proof paper and is not intended to show the paper or the ink used on the finished job.

**PLEASE READ PROOF CAREFULLY FOR ERRORS**

If proof is correct, check "OK to Print" box, sign and date it. If changes are needed, show such changes clearly in the margin of the proof, check "Send Another Proof" box, sign and date it. We cannot assume responsibility for corrections other than those indicated on proof.

If the copy as originally submitted is changed when the proof is returned, the changes in the copy are considered alterations. Alterations will be an additional charge.

**PLEASE RETURN COPY WITH PROOF**



**CORPORATE HEADQUARTERS/  
FOLDING CARTON PLANT**

28355 North Bradley Road, Lake Forest, Illinois 60045  
Tel: (847) 367-5990 Fax: (847) 367-4403

**FLEXOGRAPHIC PRINTING PLANT**

9949 58th Place, Kenosha, Wisconsin 53144  
Tel: (262) 925-4740 Fax: (262) 925-4749

Form 131R3

Operator Init. \_\_\_\_\_ Date \_\_\_\_\_

Proofer 1 Init. \_\_\_\_\_ Date \_\_\_\_\_

Proofer 2 Init. \_\_\_\_\_ Date \_\_\_\_\_

JOB HISTORY:

**PROOF A** Corrections -

**BLACK**

